

The Birkhoff theorem for topologically massive gravity

Marco Cavaglia*

Max-Planck-Institut für Gravitationsphysik

Albert-Einstein-Institut

Am Mühlenberg 5

D-14476 Golm, Germany

February 7, 2008

Abstract

We derive the general $\Sigma_2 \times S$ solution of topologically massive gravity in vacuum and in presence of a cosmological constant. The field equations reduce to three-dimensional Einstein equations and the solution has constant Ricci tensor. We briefly discuss the emergence of non-Ricci flat solutions when spin is introduced.

Pac(s) numbers: 04.50.+h, 04.20.Cv, 04.20.Jb

Keyword(s): Alternative theories of gravity, Fundamental Problems, Exact solutions

*E-mail: cavaglia@aei-potsdam.mpg.de; web page: <http://www.aei-potsdam.mpg.de/~cavaglia>

Three-dimensional gravity allows for the construction of a Chern-Simons-like term closely related to the four-dimensional Pontryagin invariant [1]. Adding this Chern-Simons term to the action and varying the latter w.r.t. the metric, one obtains field equations corresponding to Einstein equations plus a term proportional to a symmetric, traceless, covariantly constant tensor of third derivative order (Cotton tensor, $C_{\mu\nu}$) [1]

$$G^\mu{}_\nu + \frac{1}{m} C^\mu{}_\nu = \lambda \delta^\mu{}_\nu, \quad (1)$$

$$C^\mu{}_\nu = \epsilon^{\mu\sigma\kappa} \nabla_\sigma \left(R_{\nu\kappa} - \frac{1}{4} g_{\nu\kappa} R \right), \quad (2)$$

where ϵ is the completely antisymmetric Levi Civita tensor in three-dimensions. The parameter m in Eq. (1) has mass dimension (in Planck units) and we have allowed for the existence of a cosmological constant λ . This new theory is called topologically massive gravity (TMG) and has several surprising properties [1]. In particular, it has no unitarity or ghost problems and appears to be renormalizable.

A quite large amount of attention has been devoted to the investigation of the fundamental properties of TMG. In particular, TMG allows for non-trivial (non Ricci flat) solutions in three dimensions [2]. In this context, the study of exact solutions of TMG plays a major role. (Quoting Deser [3]: “[...] no-one has succeeded as yet in finding the simplest possible, ‘Schwarzschild’ solution to the nonlinear model, *i.e.*, a circularly symmetric time-independent (we don’t know if there’s a Birkhoff theorem) exterior geometry that obeys the [Eq. (1)].”)

Having this in mind, the aim of this paper is the investigation of cylindrically symmetric solutions of TMG. Given the

- **Definition 1:** A three-dimensional metric with topology $\Sigma_2 \times S$

$$ds^2 = \gamma_{\mu\nu}(x) dx^\mu dx^\nu - \rho(x)^2 d\phi^2, \quad (3)$$

where $\phi \in [0, 2\pi[$ and $\gamma_{\mu\nu}$ ($\mu, \nu = 0, 1$) is a generic hyperbolic two-dimensional metric with signature $(1, -1)$, is *static* when it possesses a Killing vector $\xi = \xi^\mu(x) \partial_\mu$ (for the definition of staticity in two-dimensional dilaton gravity see [4]);

- **Definition 2:** A solution of TMG equations is called *locally Einstein trivial* if the Cotton tensor vanishes identically when is evaluated on the solution;

we prove the following

- **Theorem:** The most general solution of Eq. (1) with topology $\Sigma_2 \times S$ is static and locally Einstein trivial.

- **Corollary:** When spin is introduced, non-static and non-Einstein trivial solutions do appear.

The above theorem can be seen as the Birkhoff theorem for TMG. As a consequence of the theorem, the most general solution of TMG with topology $\Sigma_2 \times S$ is locally flat for $\lambda = 0$ and locally De Sitter/Anti-de Sitter for $\lambda \neq 0$. As a result, TMG allows for the existence of BTZ black holes [5].

Conventions: Throughout the paper we use the signature $(+, -, -)$ and Landau-Lifshits [6] conventions for the Ricci tensor.

- **Proof:**

Let us consider the metric (3). Since in two-dimensions any metric is locally conformally flat we choose conformal light-cone coordinates u, v on Σ_2 . Equation (3) is cast in the form

$$ds^2 = f(u, v)dudv - \rho(u, v)^2 d\phi^2, \quad f(u, v) \geq 0. \quad (4)$$

Using (4) the Einstein tensor $G^\mu{}_\nu$ and the Cotton tensor read

$$G^\mu{}_\nu = 2 \begin{pmatrix} \mathcal{H}_4 & -\mathcal{H}_1/f & 0 \\ -\mathcal{H}_2/f & \mathcal{H}_4 & 0 \\ 0 & 0 & \mathcal{H}_3 \end{pmatrix}, \quad (5)$$

and

$$C^\mu{}_\nu = \begin{pmatrix} 0 & 0 & 2\rho \frac{2\partial_u \mathcal{H}_1 - f\partial_v \mathcal{H}_3}{f^2} \\ 0 & 0 & -2\rho \frac{2\partial_v \mathcal{H}_2 - f\partial_u \mathcal{H}_3}{f^2} \\ \frac{2\partial_v \mathcal{H}_2 - f\partial_u \mathcal{H}_3}{f\rho} & -\frac{2\partial_u \mathcal{H}_1 - f\partial_v \mathcal{H}_3}{f\rho} & 0 \end{pmatrix}, \quad (6)$$

respectively, where

$$\mathcal{H}_1 = \frac{f}{\rho} \partial_v \left(\frac{\partial_v \rho}{f} \right), \quad \mathcal{H}_2 = \frac{f}{\rho} \partial_u \left(\frac{\partial_u \rho}{f} \right), \quad (7)$$

$$(8)$$

$$\mathcal{H}_3 = \frac{\partial_u \partial_v (\ln f)}{f}, \quad \mathcal{H}_4 = \frac{\partial_u \partial_v \rho}{f\rho}. \quad (9)$$

(Note that $\det(C^\mu{}_\nu) = 0$.) From (1) and (5)-(6) it follows immediately

$$2\partial_u \mathcal{H}_1 - f\partial_v \mathcal{H}_3 = 0, \quad (10)$$

$$2\partial_v \mathcal{H}_2 - f\partial_u \mathcal{H}_3 = 0. \quad (11)$$

Thus the Cotton tensor vanishes and the space time is trivial. Equations (1) can be written in the form

$$\mathcal{H}_1 = \mathcal{H}_2 = 0, \quad (12)$$

$$\mathcal{H}_3 = \mathcal{H}_4 = \lambda/2. \quad (13)$$

The general solution of Eqs. (12)-(13) is

$$f(u, v) = \frac{d\rho}{dh} \frac{dU}{du} \frac{dV}{dv}, \quad \rho(u, v) = \rho(h), \quad (14)$$

where $h(u, v) \equiv U(u) + V(v)$, $U(u)$, $V(v)$ are arbitrary functions. Clearly, the solution possesses the Killing vector

$$\xi = \left(\frac{dU}{du}\right)^{-1} \partial_u - \left(\frac{dV}{dv}\right)^{-1} \partial_v. \quad (15)$$

(Actually, in the case under consideration the space time is maximally symmetric and possesses six Killing vectors.) The conformal factor ρ satisfies the ordinary differential equation

$$\frac{d\rho}{dh} = C + \frac{\lambda}{4}\rho^2, \quad (16)$$

where C is an integration constant. This concludes the proof of the theorem.

For sake of completeness, solving (16) we have ($C \neq 0$)

$$ds^2 = \frac{C}{\cos^2 \left[\frac{\sqrt{\lambda C}}{2}(U + V) + K \right]} dU dV - \frac{4C}{\lambda} \tan^2 \left[\frac{\sqrt{\lambda C}}{2}(U + V) + K \right] d\phi^2, \quad (17)$$

$$ds^2 = \frac{C}{\cosh^2 \left[\frac{\sqrt{-\lambda C}}{2}(U + V) + K \right]} dU dV + \frac{4C}{\lambda} \tanh^2 \left[\frac{\sqrt{-\lambda C}}{2}(U + V) + K \right] d\phi^2 \quad (18)$$

for $C\lambda > 0$ and $C\lambda < 0$ respectively. For $\lambda = 0$ the spacetime is locally flat, as we do expect from (14) and (16). Finally, when $C = 0$ ($\lambda \neq 0$) we obtain

$$ds^2 = \frac{1}{\left[K - \frac{\lambda}{4}(U + V) \right]^2} \left[\frac{\lambda}{4} dU dV - d\phi^2 \right], \quad (19)$$

that corresponds to De Sitter/anti-De Sitter space time in conformal coordinates and is related to (17)-(18) by a coordinate transformation.

The theorem can be (partially) extended to some matter-coupled models of TMG. For instance, the Cotton tensor must necessarily vanish when a scalar field is coupled to the system since the (u, ϕ) and (v, ϕ) components of the stress-energy tensor $T^\mu{}_\nu$ are identically zero. In this case, of course, Einstein triviality does imply neither staticity nor Ricci-flatness. The solutions of TMG are the subset of the solutions of Einstein equations that satisfy Eqs. (10)-(11).

As it has been suggested by Aliev and Nutku [7], the theorem is violated if spin is added to the model. We shall prove this by finding a counterexample. The most general three-dimensional metric with a Killing vector ∂_ϕ can be locally cast in the form

$$ds^2 = F(u, v)^2 dudv + H(u, v)dud\phi + H(u, v)dvd\phi + E(u, v)d\phi^2. \quad (20)$$

By a simple change of coordinates it is straightforward to see that the ansatz (20) describes a stationary (spinning) geometry. Indeed, setting $u = t + x$ and $v = t - x$ Eq. (20) is cast in the form

$$ds^2 = F(x, t)^2 \left[(dt + w(x, t)d\phi)^2 - dx^2 \right] - [F(x, t)^2 w(x, t)^2 - E(u, v)]d\phi^2, \quad (21)$$

where $w(x, t) = H(x, t)/F(x, t)^2$.

Let us consider for simplicity $E(u, v) = 0$ and no cosmological constant. Using Eq. (20) we find that the only non-vanishing component of the Cotton tensor is $C^{\phi\phi}$. As a consequence, the (μ, ν) ($\mu, \nu = u, v$) components of Eq. (1) (with upper indices) coincide with three-dimensional vacuum Einstein equations. They reduce to

$$(\partial_- H(u, v))^2 = 0, \quad (22)$$

where $\partial_- = \partial_u - \partial_v$. The general solution of Eq. (22) is

$$H(u, v) \equiv H(u + v). \quad (23)$$

Substituting Eq. (23) in the field equations the (μ, ϕ) ($\mu = u, v$) components of Eq. (1) are identically satisfied. The (ϕ, ϕ) component of the Cotton tensor is

$$C^{\phi\phi} = \frac{H}{|HF|} \partial_- G^{\phi\phi}, \quad (24)$$

where

$$G^{\phi\phi} = -\frac{1}{H^3 F} \left(4H \partial_{uv}^2 F - \partial_+ H \partial_+ F \right), \quad (25)$$

is the $(\phi\phi)$ component of the Einstein tensor and $\partial_+ = \partial_u + \partial_v$.

From Eq. (24) it is straightforward to prove that if the space time is static ($F \equiv F(u+v)$) it is also locally Einstein trivial because $C^{\phi\phi} = 0$. However, non-static solutions do appear. Let us choose $F \equiv F(u - v)$. Equation (25) becomes

$$2F'''F - F'' \left(2F' - m \frac{H}{|HF|} F^3 \right) = 0, \quad (26)$$

where ' means differentiation w.r.t. $u - v$. Let us consider for simplicity $H > 0$. Integrating (26) we find

$$2y'' + myy' = Ay, \quad y = |F|, \quad (27)$$

where A is an integration constant. When $A = 0$ (27) can be integrated explicitly. We obtain three different solutions

$$y_+ = \frac{\alpha}{m} \tanh \left[\frac{\alpha}{4}(u - v) - \beta \right], \quad (28)$$

$$y_- = -\frac{\alpha}{m} \tan \left[\frac{\alpha}{4}(u - v) - \beta \right], \quad (29)$$

$$y_0 = \frac{\alpha/m}{\alpha(u - v)/4 - \beta}. \quad (30)$$

The three-dimensional line elements are

$$ds_i^2 = y_i(u - v)^2 dudv + H(u + v)dud\phi + H(u + v)dvd\phi. \quad (31)$$

Note that ds_+^2 and ds_0^2 are defined in the region $\alpha(u - v)/4 - \beta > 0$ and ds_-^2 is defined in the regions $\pi(2n - 1)/2 < \alpha(u - v)/4 - \beta < n\pi$, where n is a integer number. (The regions of definition are reversed when $H(u, v) < 0$.)

Clearly, Eqs. (31) describe a non-trivial non-static geometry. Thus we have shown by an explicit example that non-static solutions do occur in spinning cylindrically symmetric TMG geometries. The presence of the Killing vector associated to cylindrical symmetry does not forbid the existence of non-static solutions.

Acknowledgments

This work was supported by a Human Capital and Mobility grant of the European Union, contract number ERBFMRX-CT96-0012. We are grateful to V. de Alfaro and Y. Nutku for useful discussions connected to the subject of this paper.

References

- [1] S. Deser, R. Jackiw, and S. Templeton, *Phys. Rev.* **D 48**, 975 (1982); *Ann. Phys.* **140**, 372 (1982).
- [2] I. Vuorio, *Phys. Lett.* **B 163**, 91 (1985);
R. Percacci, P. Sodano, I. Vuorio, *Ann. Phys.* **176**, 344 (1987);
Y. Nutku, *Ann. Phys.* **195**, 16 (1989);
M.E. Ortiz, *Class. Quant. Grav.* **7**, 1835 (1990); *Ann. Phys.* **200**, 345 (1990);

- G. Clément, *Class. Quant. Grav.* **7**, L93 (1990); **9**, 2615 (1992); *ibid.* 2635 (1992)
 B. Linet, *Gen. Rel. Grav.* **23**, 15 (1991);
 Y. Nutku, *Class. Quant. Grav.* **10**, 2657 (1993).
- [3] S. Deser, “Dimensionally Challenged Gravities”, Contribution to J. Stachel Festschrift, *preprint gr-qc/9812013*, to appear in *Boston studies in the philosophy of science*.
- [4] A.T. Filippov, *Mod. Phys. Lett.* **11**, 1691 (1996); *Int. Jou. Mod. Phys.* **A 12**, 13 (1997);
 M. Cavaglià, *Phys. Rev. D* **57**, 5295 (1998);
 T. Klösch and T. Strobl, *Class. Quantum Grav.* **13**, 965 (1996); **13**, 2395 (1997); **14**, 1689 (1997).
- [5] M. Bañados, C. Teitelboim, and J. Zanelli *Phys. Rev. Lett.* **69**, 1849 (1992).
- [6] L.D. Landau and E.M. Lifshitz, “The Classical Theory of Fields”, Pergamon Press, 1962.
- [7] A.N. Aliev, Y. Nutku, *Class. Quant. Grav.* **13**, L29 (1996).